

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12N 15/53, 15/82, 9/02, C12Q 1/26, G01N 33/53, A61K 38/44, C07K 16/40, C12P 7/64, C12N 5/10, A01H 5/00, C12Q 1/68</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/27111 (43) International Publication Date: 3 June 1999 (03.06.99)</p>
<p>(21) International Application Number: PCT/GB98/03507 (22) International Filing Date: 24 November 1998 (24.11.98) (30) Priority Data: 9724783.7 24 November 1997 (24.11.97) GB (71) Applicant (for all designated States except US): UNIVERSITY OF BRISTOL [GB/GB]; Senate House, Tyndall Avenue, Bristol BS8 1TH (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): NAPIER, Johnathan, A. [GB/GB]; I ACR-Long Ashton Research Station, Dept. of Agriculture Sciences, University of Bristol, Long Ashton, Bristol BS41 9AF (GB). (74) Agents: DEAN, John, Paul et al.; Withers & Rogers, 4 Dyer's Buildings, Holborn, London EC1N 2QP (GB).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: DESATURASE GENES AND THEIR USE</p>		
<p>(57) Abstract</p> <p>cDNA encoding <i>C. elegans</i> Δ^6 desaturase has been cloned and sequenced, and the Δ^6 desaturase amino acid sequence has been determined. The <i>C. elegans</i> Δ^6 desaturase has a surprisingly low level of sequence identity with the known borage Δ^6 desaturase. The <i>C. elegans</i> Δ^6 desaturase has been expressed in yeast. It and other desaturases can be cloned in host organisms (e.g. plants) and can be used to provide useful metabolites.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

DESATURASE GENES AND THEIR USE

The present invention relates, *inter alia*, to novel desaturases and to uses thereof.

Over the last few years a number of microsomal and soluble fatty acid desaturases have been isolated from higher plants, most notably from *Arabidopsis thaliana*. This has resulted from a combined genetic and biochemical approach to the generation and complementation of mutant *Arabidopsis* lines defective in fatty acid desaturation or elongation (Somerville C, Browse J (1996) *Trends Cell Biol.* 6, 148-1153). The importance of this approach has been validated by the isolation and characterisation of genes encoding microsomal desaturases such the Δ^{12} (Okuley J, *et al*, (1994), *Plant Cell* 6, 147-158) and Δ^{15} (Arondel V, *et al*, (1992) *Science* 258, 1353-1355) desaturases (encoded by the FAD2 and FAD3 genes respectively), enzymes which had previously proved intractable to classical purification techniques on account of their hydrophobicity. The isolation of these and related enzymes, such as the Δ^{12} hydroxylase from *Ricinus communis* (van de Loo FN *et al* (1995) *Proc. Natl. Acad. Sci USA* 92, 6743-6747), has allowed the identification of a number of conserved motifs in plant microsomal desaturases, most notably the so-called "histidine boxes" (Shanklin, J *et al* (1997) *Proc Natl. Acad. Sci USA.* 92, 6743-6747). Proteins containing these motifs can be classified as di-iron centre-containing enzymes (Shanklin, J *et al* (1997) *Proc. Natl, Acad Sci. USA* 94, 2981-1986).

WO93/11245 (Du Pont) discloses various nucleic acid fragments encoding desaturases, particularly Δ^{12} and Δ^{15} desaturases, which have been isolated from various plants. Recently a cDNA clone was isolated from the plant borage, (*Borago officinalis*) which accumulates γ -linoleic acid (GLA), using highly degenerate PCR against these histidine motifs. US5614393 (Rhone-Poulenc Agrochimie) discloses and claims the nucleotide sequence of borage Δ^6 desaturase. Whilst the specification suggests that Δ^6 desaturase-encoding nucleic acids might be isolated from animal cells without difficulty by the skilled person no suitable animal cells are suggested (in contrast to suggested fungal

and bacterial cells) and there is no disclosure of the isolation of such nucleic acids from animal cells. The isolated DNA clone was shown by heterologous expression in transgenic tobacco to encode a microsomal Δ^6 desaturase (Sayanova O *et al* (1997) *Proc. Natl. Acad. Sci. USA.* **94**, 4211-4216). Desaturation at the Δ^6 position is an unusual modification in higher plants, occurring only in a small number of species such as borage, evening primrose (*Oenothera spp.*) and redcurrant (*Ribes spp.*), which accumulate the Δ^6 -unsaturated fatty acids GLA and octadecatetraenoic acid (OTA:18:^{4,6,9,12,15}, also known as stearidonic acid) in the seeds and/or leaves.

GLA is a high value plant fatty acid, and is widely used in the treatment of a number of medical conditions, including eczema and mastalgia. It has been postulated that the application of GLA replaces the loss of, or meets an increased requirement for, endogenous Δ^6 -unsaturated fatty acids (Horrobin, D.F. (1990) *Rev. Contemp. Pharmacother.* **1**: 1-45).

For reference purposes Figure 5 is provided to show in simplified form a metabolic pathway believed to occur in certain organisms (including humans) and involving Δ^6 desaturases. It can be seen that GLA can be synthesised *in vivo* from linoleic acid under the action of a Δ^6 desaturase and that GLA can be used to synthesise dihomo-GLA, which can be converted to arachidonic acid under the influence of a Δ^5 desaturase. Arachidonic acid is a precursor of various important eicosanoids (including prostaglandins and leucotrienes). Δ^6 desaturase also converts α linoleic acid into OTA. Thus it is clear that the Δ^6 desaturase is the first committed step on the biosynthetic pathway of these biologically active molecules (see Fig. 5).

The sequence of the previously isolated borage microsomal Δ^6 desaturase differs from previously characterised plant microsomal desaturases/hydroxylases in that it contains an N-terminal extension which shows homology to cytochrome b₅, and also in that the third (most C-terminal) histidine box varies from the consensus (Shanklin J *et al* (1997) *Proc. Natl. Acad. Sci. USA* **94**, 2981-1986) H-X-X-H-H, with a glutamine replacing the first histidine. This was also observed in the case of the cyanobacteria *Synechocystis* Δ^6 desaturase (GenBank ID; L11421). WO93/06712 (Rhone Poulenc Agrochimie) discloses

an isolated nucleic acid encoding a Δ^6 desaturase isolated from the *Synechocystis*, and claims bacterial Δ^6 desaturases and their uses.

Although Δ^6 fatty acid desaturation is an unusual modification in higher plants, it is believed to be common in animals. The essential fatty acid linoleic acid (18:2 $\Delta^{9,12}$) is desaturated to GLA by a Δ^6 desaturase as a first step in the biosynthetic pathway of the eicosanoids (which include prostaglandins and leucotrienes). This results in the rapid metabolism of GLA (to di-homo-GLA and arachidonic acid; i.e. 20:3 $\Delta^{8,11,14}$ and 20:4 $\Delta^{5,8,11,14}$ respectively). Accumulation of GLA is therefore not usually observed.

The nematode worm *Caenorhabditis elegans* is extremely useful in that it has well understood genetics and has many similarities with higher animals such as humans and is therefore extremely useful in the development of desaturases for use in such animals.

According to the present invention, there is provided a polypeptide having desaturase activity, which comprises the amino acid sequence shown in Figure 1.

The amino acid sequence shown in Figure 1 is that of a Δ^6 desaturase that is present in the nematode worm *Caenorhabditis elegans*. This is highly significant since prior to the present invention no successful sequencing or purification of an animal Δ^6 desaturase had been reported. As *C. elegans* does not accumulate GLA isolation of a Δ^6 desaturase from it was an unexpected target for isolating desaturases gene in.

The desaturase of the invention is significantly different from known desaturases. The homology between the Δ^6 desaturase of the invention and the microsomal Δ^{12} and Δ^{15} desaturases from Arabidopsis described in WO93/11245 are 24% and 16% respectively as determined using the BESTFIT program. The Δ^6 desaturase gene of the present invention shows 21% identity with the *C.elegans* FAT-1 desaturase described in Spychalla, J. P. *et al* Proc. Natl Acad. Sci **94** 1142-1147 paper. The sequence homology between the Δ^6 desaturase of the present invention and the *Synechocystis* Δ^6 described in WO93/06712 is only 23%.

According to another aspect of the invention there is provided therefore an isolated animal Δ^6 desaturase.

The amino acid sequence shown in Figure 1 is also of significance because it has a very low level of sequence identity with the borage Δ^6 desaturase (the only other eukaryotic Δ^6 desaturase to have been sequenced prior to the present invention). Indeed, this level of sequence identity is below 32 %. At such a low level of identity it might be expected that the two polypeptides would have completely different functions. Unexpectedly, both have Δ^6 desaturase activity.

The present invention is, however, not limited to a Δ^6 desaturase having the sequence shown in Figure 1. It also includes other desaturases having at least 32% sequence identity therewith. Preferred polypeptides of the present invention have at least 40 % or more preferably at least 50% amino acid sequence identity therewith. More preferably the degree of sequence identity is at least 75%. Sequence identities of at least 90%, at least 95% or at least 99% are most preferred.

For the purposes of the present invention, sequence identity (whether amino acid or nucleotide) can be determined by using the "BESTFIT" program of the Wisconsin Sequence Analysis Package GCG 8.0.

Where high degrees of sequence identity are present there may be relatively few differences in amino acid sequence. Thus for example there may be less than 20, less than 10, or even less than 5 differences.

Fragments of the polypeptides described above are also within the scope of the present invention, provided that they have desaturase activity, that is to say they have the ability to introduce a double bond into a substrate at a specific position as determined by GCMS. What is the lowest limit for activity. These fragments are preferably at least 100 amino acids long. More preferably, the fragments are at least 150 amino acids long.

In summary, a polypeptide of the present invention has desaturase activity and:

- a) comprises the amino acid sequence shown in Figure 1;

- b) has one or more amino acid deletions, insertions or substitutions relative to a polypeptide as defined in a) above, but has at least 32% amino acid sequence identity therewith; or
- c) is a fragment of a polypeptide as defined in a) or b) above, which is at least 100 amino acids long.

The term "polypeptide" is used herein in a broad sense to indicate that a particular molecule comprises a plurality of amino acids joined together by peptide bonds. It therefore includes within its scope substances, which may sometimes be referred to in the literature as peptides, polypeptides or proteins.

Desirably a polypeptide of the present invention will have a cytochrome domain. A cytochrome domain can be defined as an electron-transporting domain that contains a heme prosthetic group. Preferably a cytochrome b domain is present. More preferably a cytochrome b₅ domain is present (desirably this includes a H-P-G-G-X₁₅-F-X_{3,6}-H, where X is any amino acid, motif). A cytochrome b₅ domain is present in both the borage Δ^6 desaturase and in the *C. elegans* Δ^6 desaturase amino acid sequence shown in Figure 2B. The cytochrome b₅ domain is preferably an N-terminal domain – i.e. it is closer to the N-terminal end of the desaturase than to the C-terminal end. This contrasts with other desaturases. For example, yeast Δ^9 desaturase, has a C-terminal cytochrome b₅ domain and plant Δ^{12} and Δ^{15} desaturases which do not have any b₅ domain.

A polypeptide of the present invention preferably has one or more (most preferably three) histidine boxes. One of these may have an H→Q substitution. (This provides a variant histidine box that is believed to be conserved over a range of animal / plant species.)

Polypeptides of the present invention can have any regiospecificity including *cis/trans* activity although it is preferred that they are front end desaturases that introduce a double bond between the C3 and C7 positions, measured from the COOH (Δ end) of the group. A skilled person is readily able to distinguish between different desaturases by determining the different positions of double bonds introduced by the desaturases. This can be done by known analytical techniques e.g. by using gas chromatography and mass spectrometry.

Particularly preferred desaturases of the invention are Δ^6 desaturases.

Desirably the desaturases occur naturally in one or more organisms that do not accumulate GLA (i.e. where GLA may be produced, but is not normally detectable because it is very quickly metabolised). Such desaturases may occur naturally in one or more animals. The desaturases occur naturally in one or more nematodes, e.g. in *C. elegans*.

In order to appreciate the scope of the present invention more fully, polypeptides within the scope of each of a), b) and c) above will now be discussed in greater detail.

Polypeptides within the scope of a)

A polypeptide within the scope of a) may consist of the amino acid sequence shown in Figure 1 or may have an additional N-terminal and/or an additional C-terminal amino acid sequence.

Additional N-terminal or C-terminal sequences may be provided for various reasons and techniques for providing such additional sequences are well known in the art. Such techniques include using gene-cloning techniques whereby nucleic acid molecules are ligated together and are then used to express a polypeptide in an appropriate host.

Additional sequences may be provided in order to alter the characteristics of a particular polypeptide. This can be useful in improving expression or regulation of expression in particular expression systems. For example, an additional sequence may provide some protection against proteolytic cleavage.

Additional sequences can also be useful in altering the properties of a polypeptide to aid in identification or purification. For example, a signal sequence may be present to direct the transport of the polypeptide to a particular location within a cell or to export the polypeptide from the cell. Different signal sequences can be used for different expression systems.

Another example of the provision of an additional sequence is where a polypeptide is linked to a moiety capable of being isolated by affinity chromatography. The moiety may be an epitope and the affinity column may comprise immobilised antibodies or immobilised antibody fragments that bind to said epitope (desirably with a high degree of specificity). The

resultant fusion protein can usually be eluted from the column by addition of an appropriate buffer.

Additional N-terminal or C-terminal sequences may, however, be present simply as a result of a particular technique used to obtain a polypeptide of the present invention and need not provide any particular advantageous characteristic.

Polypeptides within the scope of b)

Turning now to the polypeptides defined in b) above, it will be appreciated that these are variants of the polypeptides given in a) above.

Various changes can often be made to the amino acid sequence of a polypeptide which has a desired property in order to produce variants which still have that property. Such variants of the polypeptides described in a) above are within the scope of the present invention and are discussed in greater detail in sections (i) to (iii) below. They include allelic and non-allelic variants.

(i) Substitutions

An example of a variant of the present invention is a polypeptide as defined in a) above, apart from the substitution of one or more amino acids with one or more other amino acids.

The skilled person is aware that various amino acids have similar characteristics. One or more such amino acids of a polypeptide can often be substituted by one or more other such amino acids without eliminating a desired property of that polypeptide (such as desaturase activity).

For example, the amino acids glycine, alanine, valine, leucine and isoleucine can often be substituted for one another (amino acids having aliphatic side chains). Of these possible substitutions it is preferred that glycine and alanine are used to substitute for one another (since they have relatively short side chains) and that valine, leucine and isoleucine are used to substitute for one another (since they have larger aliphatic side chains which are hydrophobic). Other amino acids that can often be substituted for one another include phenylalanine, tyrosine and tryptophan (amino acids having aromatic side chains); lysine, arginine and histidine (amino acids having basic side chains); aspartate and glutamate (amino

acids having acidic side chains); asparagine and glutamine (amino acids having amide side chains); and cysteine and methionine (amino acids having sulphur containing side chains).

Substitutions of this nature are often referred to as "conservative" or "semi-conservative" amino acid substitutions.

(ii) Deletions

Amino acid deletions can be advantageous since the overall length and the molecular weight of a polypeptide can be reduced whilst still retaining a desired activity. This can enable the amount of polypeptide required for a particular purpose to be reduced.

(iii) Insertions

Amino acid insertions relative to a polypeptide as defined in a) above can also be made. This may be done to alter the nature of the polypeptide (e.g. to assist in identification, purification or expression).

Polypeptides incorporating amino acid changes (whether substitutions, deletions or insertions) relative to the sequence of a polypeptide as defined in a) above can be provided using any suitable techniques. For example, a nucleic acid sequence incorporating a desired sequence change can be provided by site-directed mutagenesis. This can then be used to allow the expression of a polypeptide having a corresponding change in its amino acid sequence.

Polypeptides within the scope of c)

As discussed *supra*, it is often advantageous to reduce the length of a polypeptide. Feature c) of the present invention therefore covers fragments of the polypeptides a) or b) above which are at least 100 amino acids long, but which do not need to be as long as the full length polypeptide shown in Figure 1. Desirably these fragments are at least 200, at least 300 or at least 400 amino acids long.

Various uses of the polypeptides of the present invention will now be described by way of example only.

Polypeptides of the present invention may be used, *inter alia*, in obtaining useful molecules. For example Δ^6 desaturases can be used in obtaining γ -linolenic acid (GLA) or in obtaining metabolites in respect of which GLA is a precursor. For example, octadecatetraenoic acid (OTA; $18:4\Delta^{6,9,12,15}$), a member of the *n*-3 (or ω -3) fatty acids may be produced by the Δ^6 -desaturation of α -linolenic acid.

GLA, OTA and their metabolites are useful in medicine. They can be used in the preparation of a medicament for treating a disorder involving a deficiency in GLA or of a metabolite derived *in vivo* from GLA (e.g. an eicosanoid). Disorders which may be treated include eczema, mastalgia, hypercholesterolemia, atherosclerosis, coronary disease, diabetic neuropathy, viral infections, acne, hypertension, cirrhosis and cancer.

The metabolites may be produced *in vivo* in suitable hosts or *in vitro*.

When a metabolite is to be produced *in vitro*, a desaturase of the present invention and its substrate will normally be provided separately and then combined when it is desired to produce the metabolite. The present invention therefore includes within its scope a method of making GLA or OTA comprising using a Δ^6 desaturase of the present invention to convert linoleic acid substrate or α -linolenic acid substrate to GLA or OTA respectively.

When a metabolite is to be produced *in vivo* in a organism such as a plant or animal, the substrate for a desaturase of the present invention will normally be provided by the relevant non-human organism itself. *In vivo* production of the metabolite can therefore be achieved by inserting a gene encoding a desaturase of the present invention into the organism and allowing the organism to express the desaturase. The desaturase can then act on its substrate. It will therefore be appreciated that polypeptides of the present invention can be used to provide desaturase activity in organisms that would normally not possess such activity or to increase the level of desaturase activity in organisms already having some desaturase activity. If desired, a useful metabolite may be purified from such an organism. Alternatively the organism itself may be used directly as a source of the metabolite. Particular cloning techniques that can be used to provide transgenic organisms with desaturase activity are discussed later on.

Polypeptides of the present invention can also be used as indicators of the transformation of an organism. For example, if an organism intended to be transformed does not have a particular desaturase and a nucleic acid intended for use in transformation encodes that desaturase, an assay can be performed after attempted transformation to determine whether or not the desaturase is present. Thus, in the case of the Δ^6 desaturase, an assay for the presence of GLA may be performed and GLA can serve as a simple marker for the presence of a functional transgene cassette comprising a Δ^6 desaturase encoding sequence.

A further use of the present invention is in providing antibodies. The present invention includes within its scope antibodies that bind to polypeptides of the present invention.

Preferred antibodies bind specifically to polypeptides of the present invention and can therefore be used to purify such polypeptides. (For example, they may be immobilised and used to bind to polypeptides of the present invention. The polypeptides may then be eluted by washing with a suitable eluent under appropriate conditions.)

An antibody or a derivative thereof within the scope of the present invention may be used in diagnosis. For example binding assays using such an antibody or a derivative can be used to determine whether or not a particular desaturase is present. This is useful in diagnosing disorders that arise due to the absence of the functional desaturase.

Antibodies within the scope of the present invention may be monoclonal or polyclonal.

Polyclonal antibodies can be raised by stimulating their production in a suitable animal host (e.g. a mouse, rat, guinea pig, rabbit, sheep, goat or monkey) when a polypeptide of the present invention is injected into the animal. If necessary an adjuvant may be administered together with a polypeptide of the present invention. The antibodies can then be purified by virtue of their binding to a polypeptide of the present invention.

Monoclonal antibodies can be produced from hybridomas. These can be formed by fusing myeloma cells and spleen cells which produce the desired antibody in order to form an immortal cell line. Thus the well-known Kohler & Milstein technique (*Nature* 256 52-55 (1975)) or variations upon this technique can be used.

Techniques for producing monoclonal and polyclonal antibodies that bind to a particular polypeptide are now well developed in the art. They are discussed in standard immunology textbooks, for example in Roitt *et al*, *Immunology* second edition (1989), Churchill Livingstone, London.

In addition to whole antibodies, the present invention includes derivatives thereof which are capable of binding to polypeptides of the present invention. Thus the present invention includes antibody fragments and synthetic constructs. Examples of antibody fragments and synthetic constructs are given by Dougall *et al* in *Tibtech* 12 372-379 (September 1994).

Antibody fragments include, for example, Fab, F(ab')₂ and Fv fragments. (These are discussed, for example, in Roitt *et al* (*supra*)). Fv fragments can be modified to produce a synthetic construct known as a single chain Fv (scFv) molecule. This includes a peptide linker covalently joining V_h and V_l regions, which contributes to the stability of the molecule. Other synthetic constructs that can be used include CDR peptides. These are synthetic peptides comprising antigen-binding determinants. Peptide mimetics may also be used. These molecules are usually conformationally restricted organic rings that mimic the structure of a CDR loop and that include antigen-interactive side chains.

Synthetic constructs include chimaeric molecules. Thus, for example, humanised (or primatised) antibodies or derivatives thereof are within the scope of the present invention. An example of a humanised antibody is an antibody having human framework regions, but rodent hypervariable regions.

Synthetic constructs also include molecules comprising an additional moiety which provides the molecule with some desirable property in addition to antigen binding. For example the moiety may be a label (e.g. a fluorescent or radioactive label). Alternatively, it may be a pharmaceutically active agent.

The present invention also includes nucleic acid molecules within its scope.

Such nucleic acid molecules:

- a) code for a polypeptide according to the present invention; or

- b) are complementary to molecules as defined in a) above; or
- c) hybridise to molecules as defined in a) or b) above.

These nucleic acid molecules and their uses are discussed in greater detail below:

The polypeptides of the present invention can be coded for by a large variety of nucleic acid molecules, taking into account the well-known degeneracy of the genetic code. All of these coding nucleic acid molecules are within the scope of the present invention. Preferred coding nucleic acid molecules encode the polypeptide shown in Figure 1. These include nucleic acid molecules comprising the coding sequence shown in Figure 1 and degenerate variants thereof.

The nucleic acid molecules may be used directly. Alternatively they may be inserted into vectors.

Nucleic acids or vectors containing them may be used in cloning. They may be introduced into non-human hosts to enable the expression of polypeptides of the present invention using techniques known to those skilled in the art. Alternatively, cell free expression systems may be used.

Techniques for cloning, expressing and purifying polypeptides are well known to the skilled person. Various such techniques are disclosed in standard text-books, such as in Sambrook *et al* (*Molecular Cloning* 2nd Edition, Cold Spring Harbor Laboratory Press (1989)); in Old & Primrose (*Principles of Gene Manipulation*, 5th Edition, Blackwell Scientific Publications (1994)); and in Stryer (*Biochemistry*, 4th Edition, W H Freeman and Company (1995)).

By using an appropriate expression system the polypeptides can be produced in a desired form. For example, the polypeptides can be produced by micro-organisms such as bacteria or yeast, by cultured insect cells (which may be baculovirus-infected), or by mammalian cells (such as CHO cells).

However preferred hosts are plants or plant propagating material e.g. oil seed rape, sunflower, cereals including maize, tobacco, legumes including peanut and soybean,

safflower, oil palm, coconut and other palms, cotton, sesame, mustard, linseed, castor, borage and evening primrose, or propagating material therefor.

The technology for providing plants or plant propagating material is now well developed. It is briefly discussed in WO 96/21022, for example. Desaturases isolated from animals have successfully been expressed in plants. For example, Spychalla, J.P. *et al*, (*supra*) describe the expression of a *C. elegans* desaturase in transgenic Arabidopsis. Additionally, EP0550162 (Pioneer Hi-Bred International, Inc) discloses a chimaric gene construct encoding a Δ^9 desaturase isolated from rat, and plants transformed with the construct for the production of fatty acids. The desaturase described in that publication has only 22% identity with the Δ^6 desaturase of the present invention.

Particular techniques that can be used are discussed below. It will of course be appreciated that such techniques are non-limiting.

(i) Vector systems based on Agrobacterium tumefaciens.

These include Ti based systems, such as pGV3850, in which the T-DNA has been disarmed. Desirably a selectable marker is present (e.g. a marker that provides resistance to an antibiotic).

Intermediate vectors (IVs) may also be used. They tend to be small in size and are therefore usually easier to manipulate than large Ti based vectors. IVs are generally vectors resulting from T-DNA having been cloned into *E. coli* derived plasmid vectors, such as pBR322. IVs are often conjugation-deficient and therefore a conjugation-proficient plasmid (such as pRK2013) may be used to mobilise an IV so that it can be transferred to an *Agrobacterium* recipient. *In vivo* homologous recombination can then occur in an *Agrobacterium* to allow an IV to be inserted into a resident, disarmed Ti plasmid in order that a cointegrate can be produced that is capable of replicating autonomously in the *Agrobacterium*.

Another alternative is to use binary Ti vectors. Here a modified T-DNA region carrying foreign DNA can be provided on a small plasmid that replicates in *E. coli* (e.g. pRK252). This plasmid (sometimes called mini-Ti or micro-Ti) can then be transferred conjugatively

via a tri-parental mating into an *A. tumefaciens* that contains a compatible *vir* gene (providing the *vir* function in *trans*).

Binary vectors without Ti sequences may even be used. Here bacterial *mob* and *oriT* functions may be used to promote plasmid transfer. Again, the *vir* function may be provided in *trans*.

The vector systems discussed above can be used to transfer genes into plants by using the protocol of Horsch *et al.* (*Science* 227, 1229-31 (1985)) or variants thereof. Here small discs can be punched from the leaves of a dicotyledenous plant, they can be surface-sterilised, and can then be placed in a medium including *A. tumefaciens* that contains recombinant T-DNA in which a foreign gene to be transferred is accompanied by a selectable marker (e.g. the *neo* gene). The discs can then be cultured for 2 days and then transferred to a medium for selecting the selectable marker. (This can be done for a *neo* selectable marker by culturing using a medium containing kanamycin). *A. tumefaciens* can be killed by using a carbenicillin containing medium. Shoots will normally develop from a callus after 2-4 weeks. They can then be excised and transplanted to root-inducing medium and, when large enough can be transplanted into soil.

(ii) Vector systems based on Agrobacterium rhizogenes

These include Ri derived plasmids. Ri T-DNA is generally considered not to be deleterious and therefore such plasmids can be considered as equivalent to disarmed Ti plasmids. An IV co-integrate system based on Ri plasmids has been developed.

(iii) Plant protoplast based transformation systems

Suitable techniques are described in "Plant Gene Transfer and Expression Protocols" ed. H. Jones, Human Press Methods in Molecular Biology, 49, 1995.

Transformation of plants can be facilitated by removing plant cell walls to provide protoplasts. The cell walls can be removed by any suitable means, including mechanical disruption or treatment with cellulolytic and pectinolytic enzymes. Protoplasts can then be separated from other components by centrifugation and techniques such as electroporation can then be used to transform the protoplasts with heterologous DNA. Under appropriate

culture conditions the transformed protoplasts will grow new cell walls and also divide. Shoots and roots can then be induced and plantlets formed.

(iv) Transfection by biolistics

High velocity microprojectiles carrying DNA or RNA can be used to deliver that DNA or RNA into plant cells. This has allowed a wide variety of transgenic plants to be produced and is suitable for both monocotyledenous and dicotyledenous plants. For example gold or tungsten particles coated with DNA or RNA can be used. Suitable devices for propelling the microprojectiles include gunpowder based devices, electric discharge based devices and pneumatic devices.

(v) Virus based systems

DNA plant virus vectors include cauliflower mosaic viruses (which infect a range of dicots.) and geminiviruses (which infect a wide range of dicots. and monocots). RNA plant viruses are in the majority and include Brome Mosaic Virus (which infects a number of *Graminae*, including barley) and Tobacco Mosaic Virus (which infects tobacco plants).

From the foregoing description it will be appreciated that nucleic acid molecules encoding polypeptides of the present invention can be cloned and expressed in a wide variety of organisms.

In addition to nucleic acid molecules coding for polypeptides of the present invention (referred to herein as "coding" nucleic acid molecules), the present invention also includes nucleic acid molecules complementary thereto. Thus, for example, both strands of a double stranded nucleic acid molecule are included within the scope of the present invention (whether or not they are associated with one another). Also included are mRNA molecules and complementary DNA molecules (e.g. cDNA molecules).

Nucleic acid molecules that can hybridise to one or more of the nucleic acid molecules discussed above are also covered by the present invention. Such nucleic acid molecules are referred to herein as "hybridising" nucleic acid molecules.

A hybridising nucleic acid molecule of the present invention may have a high degree of sequence identity along its length with a nucleic acid molecule within the scope of a) or b) above (e.g. at least 50%, at least 75% or at least 90% sequence identity).

As will be appreciated by those skilled in the art, the greater the degree of sequence identity that a given single stranded nucleic acid molecule has with another single stranded nucleic acid molecule, the greater the likelihood that it will hybridise to a single stranded nucleic acid molecule which is complementary to that other single stranded nucleic acid molecule under appropriate conditions.

Desirably hybridising molecules of the present invention are at least 10 nucleotides in length and preferably are at least 25, at least 50, at least 100 or at least 200 nucleotides in length.

Preferred hybridising molecules hybridise under stringent hybridisation conditions. One example of stringent hybridisation conditions is where attempted hybridisation is carried out at a temperature of from about 35°C to about 65°C using a salt solution that is about 0.9 molar. However, the skilled person will be able to vary such parameters as appropriate in order to take into account variables such as probe length, base composition, type of ions present, etc.

Most preferably, hybridising nucleic acid molecules of the present invention hybridise to a DNA molecule having the coding sequence shown in Figure 1 to an RNA equivalent thereof, or to a complementary sequence to either of the aforesaid molecules.

Hybridising nucleic acid molecules can be useful as probes or primers, for example.

Probes can be used to purify and/or to identify nucleic acids. For example they can be used to identify the presence of all or part of a desaturase gene and are therefore useful in diagnosis.

Primers are useful in amplifying nucleic acids or parts thereof, e.g. by PCR techniques.

In addition to being used as probes or primers, hybridising nucleic acid molecules of the present invention can be used as antisense molecules to alter the expression of polypeptides of the present invention by binding to complementary nucleic acid molecules. (Generally this

can be achieved by providing nucleic acid molecules that bind to RNA molecules that would normally be translated, thereby preventing translation due to the formation of duplexes.)

Hybridising molecules may also be provided as ribozymes. Ribozymes can also be used to regulate expression by binding to and cleaving RNA molecules that include particular target sequences recognised by the ribozymes.

From the foregoing discussion it will be appreciated that a large number of nucleic acids are within the scope of the present invention. Unless the context indicates otherwise, nucleic acid molecules of the present invention may therefore have one or more of the following characteristics:

- 1) They may be DNA or RNA (including variants of naturally occurring DNA or RNA structures, which have non-naturally occurring bases and/or non-naturally occurring backbones).
- 2) They may be single or double stranded.
- 3) They may be provided in recombinant form i.e. covalently linked to a heterologous 5' and/or 3' flanking sequence to provide a chimaeric molecule (e.g. a vector) which does not occur in nature.
- 4) They may be provided without 5' and/or 3' flanking sequences that normally occur in nature.
- 5) They may be provided in substantially pure form, e.g. by using probes to isolate cloned molecules having a desired target sequence or by using chemical synthesis techniques. Thus they may be provided in a form which is substantially free from contaminating proteins and/or from other nucleic acids.
- 6) They may be provided with introns (e.g. as a full-length gene) or without introns (e.g. as cDNA).

The present invention will now be described by way of example only, with reference to the accompanying drawings, Figures 1 to 6 wherein:

Fig 1 shows the DNA sequence and the deduced amino acid sequence of the full length *C. elegans* cDNA pCeD6.1. The positions of the N-terminal cytochrome b₅ domain and the variant third histidine box are underlined. The deduced amino acid sequence of this cDNA is identical to that predicted for residues 1-38 and 68-473 of W08D2.4.

Fig 2A shows a comparison of the deduced amino acid sequences of the *C. elegans* cDNA CeD6.1 and the *C. elegans* predicted protein W08D2.4. (MywormD6=CeD6.1; cew08d2=ORF W08D2.4.)

Fig 2B shows a comparison of the deduced amino acid sequences of the borage Δ^6 desaturase (Sayanova O *et al* (1997) *Proc. Natl. Acad. Sci. USA* **94**, 4211-4216) and the *C. elegans* cDNA CeD6.1. (Boofd6=*Borage officianalis* Δ^6 desaturase; ceeld6=CeD6.1.)

Fig 3 shows methyl esters of total lipids of *S. cerevisiae* grown under inducing conditions (linolate and galactose).

Fig 4 shows GC-MS analysis of the novel peak identified in yeast carrying pYCeD6.1.

Fig 5 shows a simplified version of the metabolism of n-6 essential fatty acids in mammals.

Fig 6 shows fatty acid and methyl esters of leaf material from either control transformed Arabidopsis plant (A) or transformed Arabidopsis plant expressing the *C. elegans* Δ^6 desaturase (B).

Example 1 - Isolation of Δ^6 Desaturase Gene and Expression in Yeast

The NCBI EST sequence database was searched for amino acid sequences using a known borage Δ^6 fatty acid desaturase (Sayanova O *et al* (1997) *supra*) and limiting the search to sequences containing a variant histidine box Q-X-X-H-H.

C. elegans ESTs were identified. They were further characterised by searching the *C. elegans* EST project database (Prof. Y. Kohara lab (National Institute of Genetics, Mishima, Japan); DNA Database of Japan) to identify related cosmid clones.

A partial 448 base pair cDNA clone designated as yk436b12 identified by these searches was obtained from the *C. elegans* EST project, and this was used to screen a *C. elegans* cDNA library (mixed stage; also supplied by Prof Kohara) This indicated that the clone yk436b12 was homologous to part of a gene present on cosmid W08D2 (Genbank accession number Z70271), which forms part of chromosome IV. Bases 21-2957 of cosmid W0D2 are predicted by the protein prediction program Genefinder (Wilson R *et al* (1994) *Nature* 368 32-38 to encode an ORF of 473 residues which is interrupted by 5 introns. Wilson, R. *et al* disclose part of the sequence of chromosome III of *C. elegans*. A number of positives were identified and further purified, and full length clones were confirmed by sequencing to encode a transcript likely to have been transcribed from the gene designated W08D2.4, on cosmid W08D2, as determined by database searching of the genes sequenced by the *C. elegans* genome project.

Examination of this predicted polypeptide (designated W08D2.4 by the Sanger Centre Nematode Sequencing Project, Hinxton, UK) revealed that it had a number of characteristics reminiscent of a microsomal fatty acid desaturase, including three histidine boxes. However, the predicted protein sequence indicated the presence of an N-terminal domain similar to cytochrome b_5 , containing the diagnostic H-P-G-G motif found in cytochrome b_5 proteins (Lederer F (1994) *Biochimie*. 76, 674-692). Since the Δ^6 desaturase isolated by us from borage also contained an N-terminal b_5 domain, this indicated that W0D2.4 may encode a Δ^6 desaturase.

Closer examination of the sequence revealed the presence of the variant third histidine box, with an H→Q substitution (again as observed in the borage Δ^6 desaturase). The degree of similarity between W08D2.4 and the borage Δ^6 desaturase is <52% and is therefore low. The figure of <31% obtained for identity is also low.

Since W08D2.4 was encoded by a gene containing many (6) introns, it was necessary to isolate a full length cDNA to verify the sequence predicted by the Genefinder program, and to also allow the expression of the ORF to define the encoded function.

A cDNA library was screened with the EST insert yk436b12 (generously provided by Prof Y. Kohara) and a number of positive plaques were identified. These were further purified to homogeneity, excised, and the largest inserts (of ~1450 bp) from the resulting rescued phagemids were sequenced. This confirmed that the cDNAs isolated by us were indeed homologous to W08D2.4, with the 5' and 3' ends of the cDNA being equivalent to bases 9 and 3079 of the sequence of cosmid W08D2. Since the ATG initiating codon predicted by the Genefinder program to be the start of gene product W08D2.4 was indeed the first methionine in the cDNA clone, we reasoned that we had isolated a *bona fide* full length cDNA. The DNA sequence and deduced amino acid sequence of one representative cDNA clone (termed pCeD6.1; 1463 bp in length) is shown in Fig 1; the deduced amino acid sequence is identical to that predicted for W08D2.4 over the majority of the protein. The positions of the N-terminal cytochrome b_5 domain and the variant third histidine box are underlined. The deduced amino acid sequence of this cDNA is identical to that predicted for residues 1-38 and 68-473 of W08D2.4.

However, DNA sequences encoding residues 38-67 (Y-S-I.....L-Y-F) predicted for W08D2.4 are not present in the cDNA clone. This means that the deduced amino acid sequence of CeD6.1 is in fact 443 amino acids long, as opposed to that predicted for W08D2.4, which is 473 residues in length. The only other difference between the two amino acid sequences is an M→V substitution at residue 401, resulting from an A→G base change (base 1211). The two sequences are compared in Fig 2A, as is the deduced amino acid sequence of the borage Δ^6 desaturase and that of CeD6.1 (Fig 2B). The extra sequence predicted for W08D2.4 is likely to derived from incorrect prediction of intron-exon borders.

Note the presence of the H-P-G-G cytochrome b_5 motif in the N-terminus (encoded by bases 96-108) and the H Q substitution in the third histidine box (encoded by bases 1157-1172).

The coding sequence of W08D2.4 was introduced into the yeast expression vector pYES2 by PCR. Oligonucleotides with 5' overhangs were used to introduce *KpnI* and *SacI* sites at the 5' and 3' ends respectively. The fidelity of the construct was checked by *in vitro* transcription and translation using the TnT system (Promega).

Specifically, clone pCeD6.1 was then used as a template for PCR amplification of the entire predicted coding sequence (443 amino acid residues in length), and cloned into the yeast expression vector pYES2 (Invitrogen) to yield pYCeD6. The fidelity of this PCR-generated sequence was checked *in vitro* transcription/translation of the plasmid, using the T7 RNA polymerase promoter present in pYES2.

Using the Promega TnT coupled transcription/translation system, translation products were generated and analysed by SDS-PAGE and autoradiography as per the manufacturer's instructions. This revealed (data not shown) that the plasmid pYCeD6 generated a product of ~55kD, whereas the control (pYES2) failed to yield any protein products, indicating that the construct was correct.

The resulting plasmid was introduced into yeast (*S. cerevisiae*) by the lithium acetate method (Guthrie C, Fink GR (1991) *Meths Enz* 194) and expression of the transgene was induced by the addition of galactose. The yeast was supplemented by addition of 0.2 mM linoleate (sodium salt) in the presence of 1% tergitol NP-40.

Transformation and selection of yeast able to grow on uracil-deficient medium revealed yeast colonies carrying the recombinant plasmid pYCeD6 by virtue of the URA3 selectable marker carried by pYES2. Expression of pYCeD6 was obtained by inducing the GAL promoter that is present in pYES2. This was carried out after the cells had been grown up overnight with raffinose as a carbon source, and the medium supplemented by the addition of linoleate (18:2) in the presence of low levels of detergent. This later addition was required since the normal substrate for Δ^6 desaturation is 18:2 fatty acids, which do not normally occur in *S. cerevisiae*.

Yeast total fatty acids were analyzed by GC of methyl esters. Confirmation of the presence of GLA was carried out by GC-MS (Sayanova *et al* (1997) *supra*).

In more detail, the cultures were then allowed to continue to grow after induction, with aliquots being removed for analysis by GC. When methyl esters of total fatty acids isolated from yeast carrying the plasmid pYCeD6 and grown in the presence of galactose and linoleate were analyzed by GC, an additional peak was observed (Fig 3). In Fig. 3 Panel A is yeast transformed with control (empty) vector pYES2, panel B is transformed with pYCeD6.1. The common fatty acid-methyl esters were identified as 16:0 (peak 1), 16:1 (peak 2), 18:0 (peak 3), 18:1 (peak 4), 18:2 (peak 5; supplied exogenously). The additional peak (6) in panel B corresponds to 18:3 GLA, and is indicated by an arrowhead. This had the same retention time as an authentic GLA standard, indicating that the transgenic yeast were capable of Δ^6 -desaturating linoleic acid. No such peaks were observed in any of the control samples (transformation with pYES2). The identity of this extra peak was confirmed by GC-MS, which positively identified the compound as GLA (Fig 4). In the Figure 4 experiment, the sample was analyzed for mass spectra as before (Sayanova O *et al* (1997) *Proc. Natl. Acad. Sci. USA* **94**, 4211-4216), and the data used to search a library of profiles. The sample was identified as GLA. A comparison of the mass spectra of the novel peak (A) and authentic GLA (B) is shown; visual and computer-based inspection revealed them to be identical. This confirms that CeD6.1 encodes a *C. elegans* Δ^6 desaturase, and that this cDNA is likely to be transcribed from the gene predicted to encode ORF W08D2.4, though the deduced amino acid sequence of CeD6.1 is 30 residues smaller than that of W08D2.4

Example 2 -Expression of *C.elegans* Δ^6 desaturase in plants

The coding sequence of the *C. elegans* Δ^6 desaturase was subcloned into a plant expression vector pJD330, which comprises a viral 35S promoter, and a Nos terminator. The resulting cassette or promoter/coding sequence/terminator was then subcloned into the plant binary transformation vector pBin 19, and the resulting plasmid was introduced into *Agrobacterium tumefaciens*. This *Agrobacterium* strain was then used to transform *Arabidopsis* by the vacuum-infiltration of inflorescences. Seeds were harvested and plated onto selective media containing kanamycin. Since pBin 19 confers resistance to this antibiotic, only transformed plant material will grow. Resistant lines were identified and self-fertilized to produce homozygous material. Leaf material was analyzed for fatty acid

profiles using the same method as used for the expression of the nematode desaturase in yeast. Fatty acid methyl esters were separated by GC, and novel peaks shown in Figure 6 identified by comparison with known standards and GCMS. Two novel peaks can be seen in (B) which were identified as γ -linolenic acid (peak 1) and octadecatetraenoic acid (peak 2). These are the products of Δ^6 desaturation of the precursor fatty acids linoleic acid and α -linolenic acid, respectively.

The inventors have shown that a *C. elegans* cDNA (CeD6.1) encodes a Δ^6 desaturase, and that this sequence is identical with the predicted ORF W08D2.4, except for a 30 residue insertion present in the N-terminal region of the latter protein. Whether the deduced amino acid sequence predicted for CeD6.1 represents a splicing variant of W08D2.4, or is a result of a mis-prediction of the intron/exon junctions by the Genefinder programme is unclear. However it is clear that CeD6.1 encodes a Δ^6 desaturase.

The ORF encoded by this *C. elegans* sequence appears to be related to the higher plant Δ^6 fatty acid desaturase previously isolated by us (Sayanova O *et al* (1997) *supra*), in that they both contain N-terminal domains which show homology to cytochrome b_5 . Microsomal fatty acid desaturases have been demonstrated to use free microsomal cytochrome b_5 as their electron donor (Smith MA, *et al* (1990) *Biochem. J.* **272**, 23-29, Smith MA *et al* (1992) *Biochem. J.* **287**, 141-144), and the vast majority of identified sequences for these enzymes appear not to contain this additional cytochrome b_5 domain (Okuley J *et al* (1994) *Plant Cell* **6**, 147-158, Aronel V. *et al* (1992) *Science* **258**, 1353-1355 and Napier, J.A. *et al* (1997) *Biochemical J.* **328**:717-8).

Prior to the present invention only two examples of cytochrome b_5 -domain-containing desaturases were known, one being the borage Δ^6 desaturase, and the other being the yeast microsomal Δ^9 (OLE1) desaturase (Napier JA *et al* (1997) *Biochemical J.* *supra* and Mitchell AG, Martin CE (1995) *J. Biol. Chem.* **270**, 29766-29772). OLE1, however, contains a C-terminal cytochrome b_5 domain (Napier JA *et al* (1997) *Biochemical J.* *in press* and Mitchell AG, Martin CE (1995) *J. Biol. Chem.* **270**, 29766-29772). The reason for the cytochrome b_5 may be that the Δ^6 desaturase is a "front-end" desaturase. (A "front-end" desaturation can be defined as the final desaturation reaction on the fatty acid

chain, usually introducing double bonds between a pre-existing bond and the Δ -end of the carboxy group (Mitchell AG, Martin CE (1995) *J. Biol. Chem* **270**, 29766-29772 and Aitzetmuller K, Tseegsuren, N (1994) *J. Plant Physiol.* **143**, 538-543).)

In any event, it is now believed to be the case that both a variant histidine box and an N-terminal cytochrome b_5 domain are conserved in both animals and plants, as evidenced by their presence in both the borage and nematode Δ^6 desaturases.

The invention may therefore allow the identification of other Δ^6 desaturases and also other "front-end" desaturases to be identified by the presence of these motifs.

Claims

1. A polypeptide having desaturase activity, which:
 - a) has the amino acid sequence shown in Figure 1
 - b) has one or more amino acid deletions, insertions or substitutions relative to a polypeptide as defined in a) above, but has at least 32% amino acid sequence identity therewith; or
 - c) is a fragment of a polypeptide as defined in a) or b) above, which is at least 100 amino acids long.
2. A polypeptide according to claim 1, which has a cytochrome domain.
3. A polypeptide according to claim 2, which has a cytochrome b₅ domain.
4. A polypeptide according to any preceding claim, which has at least one histidine box.
5. A polypeptide according to any preceding claim, which has three histidine boxes.
6. A polypeptide according to any preceding claim, which is a front end desaturase.
7. A polypeptide according to any preceding claim, which is a Δ^6 desaturase.
8. A polypeptide according to any preceding claim, which occurs naturally in an organism that does not accumulate GLA.
9. A polypeptide according to any preceding claim, which occurs naturally in a eukaryote.
10. A polypeptide according to any preceding claim, which occurs naturally in an animal.
11. A polypeptide according to any preceding claim, which occurs naturally in a nematode.
12. A polypeptide according to any preceding claim, which occurs naturally in *C. elegans*.

13. A polypeptide according to claim 1, which consists of the amino acid sequence shown in Figure 1 or of a part thereof.
14. A polypeptide comprising a polypeptide according to any preceding claim, when covalently linked to another moiety.
15. The use of a polypeptide according to any of claims 1 to 14 in raising or selecting antibodies.
16. The use of a polypeptide according to any of claims 1 to 14 as a marker for transformation.
17. The use of a polypeptide according to claim 16 as a marker for plant transformation.
18. An antibody or a derivative thereof which binds to a polypeptide according to any of claims 1 to 14.
19. An antibody or a derivative thereof according to claim 18, for use in diagnosis.
20. A method for assessing whether or not an organism has a polypeptide according to any of claims 1 to 14, comprising determining whether or not the organism has a polypeptide that binds to an antibody or a derivative thereof according to claim 18.
21. A method according to claim 20 in which the organism is a human.
22. A method according to claim 20 or 21 preferred *in vitro*.
23. A polypeptide according to any of claims 1 to 14, for use in medicine.
24. The use of a polypeptide according to any of claims 1 to 14 in the preparation of a medicament for treating a disorder involving a deficiency in GLA in a metabolite derived *in vivo* from GLA.
25. The use of a polypeptide according to claim 23 in which the metabolite is an eicosanoid.

26. The use according to claim 23, 24, or 25 wherein the disorder is eczema, mastalgia, hypercholesterolemia, atherosclerosis, coronary disease, diabetic neuropathy, viral infections, acne, cirrhosis, hypertension and cancer.
27. A method of making GLA comprising using a polypeptide according to any one of claims 1 to 14 to convert linoleic acid to GLA
28. A method of making OTA comprising using a polypeptide according to any one of claims 1 to 14 to convert α linoleic acid to OTA.
29. A nucleic acid molecule which:
- a) codes for a polypeptide according to any of claims claim 1 to 14,
 - b) is the complement of a nucleic acid molecule as defined in a) above, or
 - c) hybridises to a nucleic acid molecule as defined in a) or b) above.
30. A vector comprising a nucleic acid molecule according to claim 29.
31. A host comprising a nucleic acid molecule according to claim 27 or a vector according to claim 30.
32. A host according to claim 31, which is a plant or plant propagating material.
33. A host according to claim 31 or claim 32, which is oil seed rape, sunflower, cereals including maize, tobacco, legumes including peanut and soybean, safflower, oil palm, coconut and other palms, cotton, sesame, mustard, linseed, castor, borage and evening primrose; or which is propagating material for any of the aforesaid.
34. A method for obtaining a polypeptide according to any of claims 1 to 14, comprising incubating a host according to any of claims 31 to 33 under conditions causing expression of said polypeptide and then purifying said polypeptide.
35. The use of nucleic acid molecule according to claim 29 as a probe or as a primer.

36. The use of a nucleic acid molecule according to claim 29 or a vector according to claim 30 for preparing an organism that accumulates GLA or a metabolite derived from GLA in that organism.
37. The use of a nucleic acid molecule according to claim 29 or a vector according to claim 30 for preparing an that is chill resistant .
38. A method of producing a host according to any of claims 31 to 33, comprising incorporating a nucleic acid according to claim 29 or a vector according to claim 30 into an organism.

1/9

10 30 50
GCTCACCAAAATGGTCGTCGACAAGAATGCCTCCGGGCTTCGAATGAAGGTCGATGGCAA
M V V D K N A S G L R M K V D G K

70 90 110
ATGGCTCTACCTTAGCGAGGAATTGGTGAAGAAACATCCAGGAGGAGCTGTTATTGAACA
W L Y L S E E L V K K H P G G A V I E Q

130 150 170
ATATAGAAATTCGGATGCTACTCATATTTTCCACGCTTTCCACGAAGGATCTTCTCAGGC
Y R N S D A T H I F H A F H E G S S Q A

190 210 230
TTATAAGCAACTTGACCTTCTGAAAAAGCAGGAGAGCACGATGAATTCCTTGAGAAACA
Y K Q L D L L K K H G E H D E F L E K Q

250 270 290
ATTGGAAGAGAGACTTGACAAAGTTGATATCAATGTATCAGCATATGATGTCAGTGTTGC
L E K R L D K V D I N V S A Y D V S V A

310 330 350
ACAAGAAAAGAAAATGGTTGAATCATTGAAAACTACGACAGAAGCTTCATGATGATGG
Q E K K M V E S F E K L R Q K L H D D G

370 390 410
ATTAATGAAAGCAAATGAAACATATTTCTGTTTAAAGCGATTTC AACACTTTCAATTAT
L M K A N E T Y F L F K A I S T L S I M

430 450 470
GGCATTGCAATTTTATCTTCAGTATCTTGGATGGTATATTACTTCTGCATGTTTATTAGC
A F A F Y L Q Y L G W Y I T S A C L L A

490 510 530
ACTTGCAATGGCAACAATTCGGATGGTTAACACATGAGTTCTGCCATCAACAGCCAACAAA
L A W Q Q F G W L T H E F C H Q Q P T K

FIG. 1

SUBSTITUTE SHEET (RULE 26)

2/9

550 570 590
GAACAGACCTTTGAATGATACTATTTCTTTGTTCTTTGGTAATTTCTTACAAGGATTTTC
N R P L N D T I S L F F G N F L Q G F S

610 630 650
AAGAGATTGGTGGAAGGACAAGCATAACACTCATCACGCTGCCACAAATGTAATTGATCA
R D W W K D K H N T H H A A T N V I D H

670 690 710
TGACGGTGATATCGACTTGGCACCACCTTTTCGCATTTATTCCAGGAGATTGTGCAAGTA
D G D I D L A P L F A F I P G D L C K Y

730 750 770
TAAGGCCAGCTTTGAAAAAGCAATTCTCAAGATTGTACCATATCAACATCTCTATTTTAC
K A S F E K A I L K I V P Y Q H L Y F T

790 810 830
CGCAATGCTTCCAATGCTCCGTTTCTCATGGACTGGTCAGTCAGTTCAATGGGTATTCAA
A M L P M L R F S W T G Q S V Q W V F K

850 870 890
AGaGAATCAAATGGAGTACAAGGTCTATCAAAGAAATGCATTCTGGGAGCAAGCAACAAT
E N Q M E Y K V Y Q R N A F W E Q A T I

910 930 950
TGTTGGACATTGGGCTTGGGTATTCTATCAATTGTTCTTATTACCAACATGGCCACTTCG
V G H W A W V F Y Q L F L L P T W P L R

970 990 1010
GGTTGCTTATTTTATTATTTTACAAATGGGAGGAGGCCTTTTGATTGCTCACGTAGTCAC
V A Y F I I S Q M G G G L L I A H V V T

FIG. 1 CONT'D

SUBSTITUTE SHEET (RULE 26)

3/9

1030 1050 1070
TTTCAACCATAACTCTGTTGATAAGTATCCAGCCAATTCTCGAATTTTAAACAACCTTCGC
F N H N S V D K Y P A N S R I L N N F A

1090 1110 1130
CGCTCTTCAAATTTTGACCACACGCAACATGACTCCATCTCCATTCATTGATTGGCTTTG
A L Q I L T T R N M T P S P F I D W L W

1150 1170 1190
GGGTGGACTCAATTATCAGATCGAGCACCCTTGTTCCTCAACAATGCCACGTTGCAATCT
G G L N Y Q I E H H L F P T M P R C N L

1210 1230 1250
GAATGCTTGCGTGAAATATGTGAAAGAATGGTGCAAAGAGAATAATCTTCCTTACCTCGT
N A C V K Y V K E W C K E N N L P Y L V

1270 1290 1310
CGATGACTACTTTGACGGATATGCAATGAATTTGCAACAATTGAAAAATATGGCTGAGCA
D D Y F D G Y A M N L Q Q L K N M A E H

1330 1350 1370
CATTCAAGCTAAAGCTGCCTAAACAATCTGGGTGTTCAAAAAGTTTTTCTTGTTTTTTT
I Q A K A A *

1390 1410 1430
AAATTTAATTCTTTGAAATTATTTGTTTTCCGTCATTCTTCCTCCATTCCCTTTTCTGGT

1450
AGAAATAAAACCTTGTTTTTCAA

FIG. 1 CONT'D

FIG. 2A

PRETTYBOX of: des.msff(*) November 4, 1997 18:33:04.76



4 / 9

Mywormd6 Cew08d2	MVVDKKNASGL MVVDKKNASGL	RMKVDGKWL RMKVDGKWL	LSEELVKKHP LSEELVKKHP	GGAVIEQ GGAVIEQ	YSI YSI	PPLNKNIE PPLNKNIE	TR TR	GIITTRGSSN GIITTRGSSN	37 60
Mywormd6 Cew08d2	YRN ALDILYFYRN	SDATHIFHAF SDATHIFHAF	HEGSSQAYKQ HEGSSQAYKQ	LDLLKKKHGEH LDLLKKKHGEH	DEFLEKQLEK DEFLEKQLEK	RLDKVDINVS RLDKVDINVS	90 120		
Mywormd6 Cew08d2	AYDVSVAAQEK AYDVSVAAQEK	KMVESFEKLR KMVESFEKLR	QKLHDDGGLMK QKLHDDGGLMK	ANETYFLFKA ANETYFLFKA	ISTLSINAF ISTLSINAF	FYLYLGMVI FYLYLGMVI	150 180		
Mywormd6 Cew08d2	TSACLLALAW TSACLLALAW	QFQGLTHER QFQGLTHER	CHQOPTKNRP CHQOPTKNRP	LNDTISLFFG LNDTISLFFG	NFLQGFSTRD NFLQGFSTRD	WKDKHNTHHA WKDKHNTHHA	210 240		
Mywormd6 Cew08d2	ATNVIDHDGD ATNVIDHDGD	IDLAPLFAFI IDLAPLFAFI	PGDLCKYKAS PGDLCKYKAS	FEKAILKIVP FEKAILKIVP	YQHLTYFTAML YQHLTYFTAML	PMLRFSWTGQ PMLRFSWTGQ	270 300		
Mywormd6 Cew08d2	SVQWVFKEHQ SVQWVFKEHQ	MEYKVYQRNA MEYKVYQRNA	FWEQATIVGH FWEQATIVGH	WAWVFYQLFL WAWVFYQLFL	LPTWPLRVAY LPTWPLRVAY	FIISQMGGL FIISQMGGL	330 360		
Mywormd6 Cew08d2	LIAHVVTENH LIAHVVTENH	NSVDKYPANS NSVDKYPANS	RILNNFAALO RILNNFAALO	ILTRNNTPS ILTRNNTPS	PFIDMLWGG PFIDMLWGG	NYQIEHHLFP NYQIEHHLFP	390 420		
Mywormd6 Cew08d2	TMPRCNLNAC TMPRCNLNAC	VKYVKEWCKE MKYVKEWCKE	NNLPYLVDV NNLPYLVDV	FDGYAHNLQ FDGYAHNLQ	LKNMAEHIOA LKNMAEHIOA	KAA*443 KAA*473			

4/9

boofd6 ceeld6	b ₅										5/9									
	MAAQIKKYYIT	SDELKKNHDKKP	GD	LWISIQGK	AYDVS	.DMVK	DHPGG	SFPLK	SLAGQEV	TDA	59									
MVDKNA	SG	LRMKVDGK	WLYLSEELVK		KHPGG	AV	.IE	QYRNSDA	THI	46								
boofd6 ceeld6	FVAFHPAS	..	TWKNL	DKF	..	FTGYYLK	DY	..	SVSE	V8	100									
	FHAFHEGSSQ	AYKQL	DL	LKK	HGEHDEFLEK	QLEKRLDKVD	INV	SA	YD	V8	106									
boofd6 ceeld6	SKMGLYDKKG	HIMFA	..	TLC	FIAMLFAMSV	YGVLF	CEGV	L	VHLF	.SG	CLM	157								
	EKLRLQKLHDD	GLMKANE	T	YF	LFKAISTLSI	MAFAFY	LQYL		GWYIT	SA	CLL	166								
boofd6 ceeld6	GHDAGHYMVV	SDSR	LN	KFMG	I	FAANCL	SGI	SIG	WWKKN	HN	217									
	THEFC	HQQPT	KNRPP	LNDTIS	L	FFGN	FLQGF	SRD	MMKDK	HN	225									
boofd6 ceeld6	LVVSSKFF	FGS	LTSHFY	EKRRL	T	EDSLSR	FFV	SYQH	WT	FYPI	277									
	...L	FAF	IPGDL	CKYKA	S	EKA	ILKIV	PYOH	LYE	TAM	279									
boofd6 ceeld6	NVSYRAHELL	G	CLVFSI	WYPL	L	LVSC	L	PNWGE	327									
	QMEYKVYQRN	AFWEQ	ATIVG	HWAW	.VFY	QL	FL	..	LPT	WPL	336									
boofd6 ceeld6	SLNHFS	SSVY	VGKPKG	NNW	FEK	QTDG	TLD	ISCP	PPWM	DWF	386									
	TFNHNS	VDKY	PANSRI	LNNF	AAL	O	IIL	TNRN	MT	PS	PFIDWL	396								
boofd6 ceeld6	LRKISPY	VIE	LCKKH	NLPYN	YASF	SKANEM	T	LR	TNR	TA	444									
	LNACVKY	VKE	WCKEN	NLPYL	VDDY	FDGYAM	NL	Q	LKN	MAE	443									
boofd6 ceeld6	HTHG	*448																		
	443																	

FIG. 2B




7/9

RESULTS OF PBM SEARCH USING THE wileybbs LIBRARY

Run =DOM10004 Scan=738 (Sub) 100%=413600 ADC Mass Range=40-456
 23 Sep 97 3:50 Compacted SLRP +EI 1UL C.E O/N INDUCTION + LA

Serial	Rel. (Sim)	Rel. (Same)	Mol. Wt.	Formula & Name
77275	99	81	292	C19 H32 O2 6, 9, 12-Octadecatrienoic acid, methyl ester
81040	95	74	122	C9 H14 1, 4-Cyclononadiene
43157	95	74	292	C19 H32 O2 6, 9, 12-Octadecatrienoic acid, methyl ester
77274	93	68	292	C19 H32 O2 6, 9, 12-Octadecatrienoic acid, methyl ester
6892	60	35	136	C10 H16 .BETA.-FENCHENE
4278	59	33	122	C9 H14 3-Nonen-1-yne, (z)-
25116	55	29	206	C15 H26 5-Pentadecen-7-yne, (z)-
17742	55	29	178	C13 H22 3-Tridecen-1-yne, (z)-
13423	55	29	162	C12 H18 1, 4, 8-Dodecatriene, (E, E, E)-
10169	34	11	150	C11 H18 Cyclopropane, 1-etheny-2-hexenyl-, 1.alpha., 2.beta. (E)
2366	33	10	108	C8 H12 1, 4-Cyclooctadiene, (z, z)-
2372	32	9	108	C8 H12 Bicyclo 5.1.0. oct-3-ene
6909	24	6	136	C10 H16 Cyclooctene, 3-ethenyl-
10171	23	6	150	C11 H18 Cyclohexene, 3-(3-methyl-1-butenyl)-, (E)-
29046	23	6	222	C15 H26 0 5, 10-Pentadecadienal, (z, z)-
17743	21	5	178	C13 H22 3-Tridecen-1-yne, (E)-
10173	20	4	150	C11 H18 Spiro 5.5.undec-1-ene
4281	20	4	122	C9 H14 Bicyclo 5.1.0. octane, 8-methylene-
10192	18	3	150	C11 H18 (-)-2-METHYL-2-BORNENE
4291	18	3	122	C9 H14 1, 2-CYCLONONADIENE
2597	18	3	110	C8 H14 Cyclopentane, (1-methylethylidene)-
7335	12	2	138	C10 H18 4-Decyne
67891	11	2	82	C6 H10 Cyclopropane, 1, 2-dimethyl-3-methylene-, cis-
618	11	2	82	C6 H10 Cyclopropane, 1, 2-dimethyl-3-methylene-, trans-
71289	4	1	136	C10 H16 2-.BETA.-PINENE

Run =DOM10004 Scan=738 (Sub) 100%=413600 ADC Mass Range=40-456
 23 Sep 97 3:50 Compacted SLRP +EI 1UL C.E O/N INDUCTION + LA



wileybbs: 77275 Rel(sim):99 Rel(same):81
 6,9,12-Octadecatrienoic acid, methyl ester
 C19 H32 O2 MW=292.240230 CAS=2676417

Me (CH₂)₄ CH CH CH₂ CH CH CH₂ CH CH (CH₂)₄ C(O) OMe




FIG. 4

SUBSTITUTE SHEET (RULE 26)




FIG. 5

9 / 9

FIG. 6A





FIG. 6B



INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03507

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/53 C12N15/82 C12N9/02 C12Q1/26 G01N33/53
A61K38/44 C07K16/40 C12P7/64 C12N5/10 A01H5/00
C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12Q G01N A61K A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HILLIER, L., ET AL.: "pk42c09.r1 Caenorhabditis briggsae cDNA similar to SP:S35157 S35157 DELTA(6)-DESATURASE - SYNECHOCYSTIS" EMBL ACCESSION NO. R05219, 18 April 1995, XP002099441 see the whole document ---	1-13, 29-31, 38
X	SWINBOURNE, J., ET AL.: "Caenorhabditis elegans cosmid W08D2" EMBL ACCESSION NO. Z70271, 23 March 1996, XP002099442 see translation product for gene W08D2.4 and sequence 1251-1999 --- -/--	1-13, 29-31, 38

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 April 1999

Date of mailing of the international search report

23/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03507

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SWINBOURNE, J., ET AL.: "T13F2.1" TREMBL ACCESSION NO. 094044, 1 February 1997, XP002099443 & EMBL ACCESSION NO. Z81122, 21 October 1996, ---	1-6, 8-13, 29-31
X	KOHARA, Y.: "C. elegans cDNA clone yk436b12 : 5' end, single read" EMBL SEQUENCE ACCESSION NO. C47394, 9 September 1997, XP002099444 see the whole document ---	29-31, 38
X	KOHARA, Y.: "C. elegans cDNA clone yk436b12 : 3' end single read" EMBL ACCESSION NO. C35646, 6 September 1997, XP002099445 see the whole document ---	29-31, 38
P, X	SAYANOVA, O., ET AL.: "Caenorhabditis elegans delta6-fatty-acid-desaturase mRNA, complete cds." EMBL ACCESSION NO. AF031477, 27 April 1998, XP002099446 see the whole document ---	1-13, 29-31, 38
P, X	-& NAPIER, J.A., ET AL.: "Identification of a Caenorhabditis elegans delta6-fatty-acid-desaturase by heterologous expression in Saccharomyces cerevisiae" BIOCHEMICAL JOURNAL, vol. 330, no. 2, March 1998, pages 611-614, XP002099453 see the whole document ---	1-13, 27, 29-31, 36, 38
P, X	WO 98 46763 A (THURMOND JENNIFER ; CALGENE LLC (US); ABBOTT LAB (US); KNUTZON DEBO) 22 October 1998 see seq IDs 2, 5, and 10. ---	1-9, 13, 23-31, 36, 38
P, X	WO 98 46764 A (THURMOND JENNIFER ; CALGENE LLC (US); ABBOTT LAB (US); KNUTZON DEBO) 22 October 1998 see sequence IDs 2 and 12 ---	1-9, 13, 23-33, 36, 38
	-/--	

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/GB 98/03507

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SAYANOVA, O., ET AL.: "Expression of a borage desaturase cDNA containing an N-terminal cytochrome b5 domain results in the accumulation of high levels of delta-6-desaturated fatty acids in transgenic tobacco"</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA.,</p> <p>vol. 94, April 1997, pages 4211-4216,</p> <p>XP002099447</p> <p>WASHINGTON US</p> <p>cited in the application</p> <p>see the whole document</p> <p style="text-align: center;">---</p>	1-38
A	<p>TANAKA, T., ET AL.: "Effects of growth temperature on the fatty acid composition of the free-living nematode <i>Caenorhabditis elegans</i>"</p> <p>LIPIDS,</p> <p>vol. 31, no. 11, - 1996 pages 1173-1178,</p> <p>XP002099448</p> <p>see page 1176, left-hand column</p> <p style="text-align: center;">---</p>	1-38
A	<p>SPYCHALLA, J. P., ET AL.: "Identification of an animal omega-3 fatty acid desaturase by heterologous gene expression in <i>Arabidopsis</i>"</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA.,</p> <p>vol. 94, February 1997, pages 1142-1147,</p> <p>XP002099449</p> <p>WASHINGTON US</p> <p>cited in the application</p> <p>see the whole document</p> <p style="text-align: center;">---</p>	1-38
A	<p>EP 0 410 637 A (IRELAND BIORESEARCH ;HEADON DENIS ROBERT (IE); MCSWEENEY FINBARR M) 30 January 1991</p> <p>see the whole document</p> <p style="text-align: center;">---</p>	15,18-22
A	<p>EP 0 454 102 A (EFAMOL HOLDINGS)</p> <p>30 October 1991</p> <p>see the whole document</p> <p style="text-align: center;">---</p>	23-26
A	<p>EP 0 400 547 A (ABBOTT LAB)</p> <p>5 December 1990</p> <p>see the whole document</p> <p style="text-align: center;">---</p>	23-26
A	<p>WO 96 21022 A (RHONE POULENC AGROCHIMIE)</p> <p>11 July 1996</p> <p>see page 2, line 3 - page 3, line 25</p> <p>see page 11, line 10 - line 21</p> <p>see page 12, line 5 - line 23</p> <p>see page 19, line 25 - page 20, line 10</p> <p>see page 31 - page 43; figures 4,9</p> <p style="text-align: center;">---</p>	24-38
	-/--	

INTERNATIONAL SEARCH REPORT

Inter. onal Application No

PCT/GB 98/03507

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>KODAMA H ET AL: "GENETIC ENHANCEMENT OF COLD TOLERANCE BY EXPRESSION OF A GENE FOR CHLOROPLAST OMEGA-3 FATTY ACID DESATURASE IN TRANSGENIC TOBACCO"</p> <p>PLANT PHYSIOLOGY, vol. 105, 1 January 1994, pages 601-605, XP002001002 see the whole document -----</p>	37

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03507

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9846763 A	22-10-1998	AU 6961698 A	11-11-1998
		AU 7114798 A	11-11-1998
		WO 9846764 A	22-10-1998
WO 9846764 A	22-10-1998	AU 7114798 A	11-11-1998
		AU 7114898 A	11-11-1998
		WO 9846765 A	22-10-1998
		AU 6961698 A	11-11-1998
		WO 9846763 A	22-10-1998
EP 0410637 A	30-01-1991	AU 5913890 A	24-01-1991
		CA 2021186 A	20-01-1991
		JP 3065652 A	20-03-1991
EP 0454102 A	30-10-1991	AT 87825 T	15-04-1993
		AU 618814 B	09-01-1992
		AU 3597489 A	14-12-1989
		AU 633442 B	28-01-1993
		CA 1334004 A	17-01-1995
		EP 0347056 A	20-12-1989
		ES 2053990 T	01-08-1994
		IE 63303 B	05-04-1995
		JP 2032017 A	01-02-1990
		JP 2796838 B	10-09-1998
		KR 129666 B	09-04-1998
		NZ 239126 A	27-07-1997
		US 4977187 A	11-12-1990
		US 5120760 A	09-06-1992
EP 0400547 A	05-12-1990	AU 633320 B	28-01-1993
		AU 5622290 A	10-01-1991
		CA 2018096 A	02-12-1990
		JP 3047119 A	28-02-1991
		US 5196198 A	23-03-1993
WO 9621022 A	11-07-1996	US 5614393 A	25-03-1997
		AU 4673596 A	24-07-1996
		BR 9510411 A	19-05-1998
		CA 2207906 A	11-07-1996
		CN 1177379 A	25-03-1998
		EP 0801680 A	22-10-1997
		JP 10511848 T	17-11-1998
		US 5789220 A	04-08-1998